

# Fludeoxyglucose ( $^{18}\text{F}$ )



[ $^{18}\text{F}$ ]Fludeoxyglucose ([ $^{18}\text{F}$ ]FDG) is the most commonly used radiopharmaceutical in clinical positron emission tomography (PET) imaging. It is a radioactive glucose analog and accumulates in tissues with high glucose metabolism. [ $^{18}\text{F}$ ]FDG is mostly used in the field of oncology and provides highly accurate diagnosis and assessment of disease stage and therapeutic response.

## Product specifications

Fludeoxyglucose ( $^{18}\text{F}$ )
185 MBq/ml at calibration time and date
Solution for injection
Store in the original package at room temperature
Expiry is 6:00 pm for calibration 10:00 am, 11:00 pm for calibration 2:00 pm
pH 4.5 – 8.5
Radiochemical purity $\geq$ 95%



✓ **AVAILABILITY:**  
Monday to Friday (Saturday on special request)

✓ **CALIBRATION:**  
10:00 am, 2:00 pm CET same day  
(5:00 pm on special request)

✓ **PACKAGING:**  
15 ml multi-dose colorless glass vial – Type I

✓ **ORDERING:**  
Curium Pharma

## Physical Data

Rad. Type	Energy (keV)	Radiation Intensity (%)
B+	249.8	96.7
E-AU-K	0,52	3,07
G-AN	511	193

## Decay Table

Physical half-life: 109.77 min

Hours\min	0	10	20	30	40	50
0	1,000	0,939	0,881	0,827	0,777	0,729
1	0,685	0,643	0,603	0,567	0,532	0,499
2	0,469	0,440	0,413	0,388	0,364	0,342
3	0,321	0,301	0,283	0,266	0,249	0,234
4	0,220	0,206	0,194	0,182	0,171	0,160
5	0,150	0,141	0,133	0,125	0,117	0,110
6	0,103	0,097	0,091	0,085	0,080	0,075

To calculate a precalibration activity, divide the activity at calibration time by the decay factor.  
For a postcalibration activity, multiply the activity at calibration time by the decay factor.

by **CYCLOTRON**<sup>VU</sup>  
radiopharmaceuticals and radionuclides

## ADDRESS

van der Boechorststraat 6A  
1081 BT Amsterdam  
The Netherlands

## CONTACT

 +31 20 44 49 123  
 info@cyclotron.nl  
 www.cyclotron.nl